

Question 1 3x10=30 points

(A) Consider the sentence $((\neg A \vee \neg B) \Rightarrow (A \Rightarrow \neg B)) \wedge (A \vee B)$

- Is this sentence satisfiable?
- Is this sentence valid?
- Is this sentence a contradiction?
- None of the above

Choose the correct answer and explain.

(B) Consider the pair of sentences $(\neg A \vee \neg B) \Rightarrow \neg(A \wedge B)$ and $((A \Rightarrow B) \vee (B \Rightarrow C))$. Are the sentences logically equivalent? Explain.

(C) Derive a weighted partial MAXSAT formula that incorporates the following statements about geese:

1. All geese are white.
2. Geese often have two legs.
3. It is very likely that a goose is either white or has two legs or both.
4. If a goose does not have wings, it cannot fly.

You may use the following variables:

W = Goose has wings

X = Goose is white

Y = Goose has two legs

Z = Goose can not fly

Question 2 2x10=20 points

(A) Translate the following sentences from Description Logic *ALC* to English language and vice versa. Suppose that the concept names are *Kid* and *Ice-Cream* and the role name is *likes*.

- $\exists \text{like}. \text{Ice} - \text{Cream} \sqsubseteq \text{Kids}$
- Every kid likes something
- Everybody likes ice-cream

(B) Use a tableau to prove whether $(\forall r. A \sqcap \forall r. B) \sqsubseteq \forall r. (A \sqcap B)$ is satisfiable or not.

Question 3 10 points

Use the Davis Putnam procedure to prove or disprove satisfiability of the formula
 $(X \vee Y \vee Z) \wedge (X \vee \neg Y) \wedge (Y \vee \neg Z) \wedge (Z \vee \neg X) \wedge (\neg X \vee \neg Y \vee \neg Z)$
 Label each step in your derivation with the part of the DP algorithm that you applied.

Question 4 2x5= 10


- Give a pseudocode description of the GSAT procedure.
- Explain the advantages and disadvantages of short and long restart intervals.

Question 5 10 points

Check out the statements below whether they are true or not, and label them accordingly with either "true" or "false".

- If $\alpha \models \beta$ and $\Pr(\alpha)=0$, then $\Pr(\beta)=0$.
- If $\alpha \models \beta \models \gamma$, then $\Pr(\alpha \mid \beta) \geq \Pr(\alpha \mid \gamma)$.


c) In the interaction graph G below, if we eliminate 4, we get G'



d) Assume G' above, according to the order elimination heuristic of "choosing the node with smallest degree", one possible elimination order is: 1, 5, 3, 2.

Question 6 2x10= 20 points

Consider the Bayesian network in the figure, and answer the questions below accordingly.



a) Are the statements below true or false? Cross the correct option:

$\text{IPr}(D, \emptyset, C)$. [True or False]

$\text{IPr}(E, \{A\}, B)$. [True or False]

$\text{IPr}(\{BD\}, \{A, C\}, E)$. [True or False]

$\text{IPr}(C, \{B, E\}, A)$. [True or False]

b) 1. Calculate the probability $\text{Pr}(A=\text{true}, B=\text{false}, E=\text{true})$.

2. Calculate the probability $\text{Pr}(A=\text{false}, B=\text{true} \mid C = \text{true})$.