

Exam: Multivariate Statistics

Code: E\_EOR2\_MS

Examinator: Andre Lucas

Co-reader: Julia Schaumburg

Date: May 20, 2019

Time: 12:00

Duration: 2 hours

Calculator allowed: Yes

Graphical calculator  
allowed: Yes

Number of questions: 4

Type of questions: Open

Answer in: English

Remarks: Take the number of points as a guidance on the maximum time to spend on that question.
---

Credit score: See the credits for each separate question

Grades: The grades will be made public latest on June 2, 2019.

Inspection: Tuesday, June 4, 2019 at 10am. Room tba via CANVAS.

Number of pages: 6 (including front page)

**Good luck!**

**Question 1. [40points]**

You have two independent random variables,  $R$  and  $A$ , with  $f_A(a) = \frac{1}{\pi}$  for  $-\frac{1}{2}\pi \leq a \leq \frac{1}{2}\pi$ , and zero else, and  $f_R(r) = \exp(-a)$  for  $a > 0$  and zero else. We therefore have that  $A$  is a point on the half-circle, and  $R$  is its ray.

A) Define  $Y_1 = R \cdot \cos(A)$  and  $Y_2 = R \cdot \sin(A)$ . Derive the multivariate cdf of  $Y = (Y_1, Y_2)^\top$ . [20pt]  
[Hint: it is convenient for deriving the expression of the inverse transformation to use the fact that  $R$  is the ray along the point of the circle (Pythagoras).]

You have a normally distributed random vector  $X = (X_1, X_2, X_3)^\top$ , with

$$X_1|X_2 \sim N(X_2, 3), \quad X_2|X_1 \sim N(1 + 0.25X_1, \star), \quad X_3| \begin{pmatrix} X_1 \\ X_2 \end{pmatrix} \sim N(X_2 - X_1, 1).$$

B) Provide the mean and covariance matrix of  $X$ . [20pt]

[Hint1: first solve for the distribution of  $(X_1, X_2)$ .]

[Hint2: if you do not succeed in solving for the distribution of  $(X_1, X_2)$ , then \*(incorrectly) assume\*  $(X_1, X_2)^\top \sim N((1, 2)^\top, \begin{pmatrix} 3 & 2 \\ 2 & 2 \end{pmatrix})$  and proceed to derive the distribution of  $X$ .]

**Question 2. [40pt]**

A) Provide the algorithm to simulate from a 4 dimensional pdf that is characterized by  $\chi^2(\nu_i)$ -marginals for  $i = 1, \dots, 4$ , and a Gaussian copula with correlation parameter  $R \in \mathbb{R}^{4 \times 4}$ . [10pt]

Let  $C^G(u_1, u_2, u_3; R)$  be the trivariate Gaussian copula with parameter  $R$ , and  $C^{St}(u_1, u_2, u_3; R, \nu)$  the Student's t copula with parameters  $R$  and  $\nu$ .

You construct a new copula as a mixture of these previous two, with mixture weights  $w = 0.2$  and  $1 - w = 0.8$ , respectively.

B) Describe the algorithm to simulate from this mixture of the two copulas. [5pt]

[Hint: just concentrate in how to draw from a mixture distribution. Skip how to draw from a copula, which was question 2A above]

C) Show whether or not the mixture of the two copulas

$C(u_1, u_2, u_3; R, \nu, w) = 0.2 \cdot C^G(u_1, u_2, u_3; R) + 0.8 \cdot C^{St}(u_1, u_2, u_3; R, \nu)$   
is itself a copula or not. [15pt]

Assume the marginal distributions

$$\begin{aligned} f_{X_1}(x_1) &= 2 \cdot \exp(-2x_1), & F_{X_1}(x_1) &= 1 - \exp(-2x_1), \\ f_{X_2}(x_2) &= 4 \cdot \exp(-4x_2), & F_{X_2}(x_2) &= 1 - \exp(-4x_2), \\ f_{X_3}(x_3) &= 3 \cdot \exp(-3x_3), & F_{X_3}(x_3) &= 1 - \exp(-3x_3), \end{aligned}$$

and assume that  $X = (X_1, X_2, X_3)^\top$  has the mixture copula dependence structure from question (C).

D) Given a realization  $X = x = (2, 2, 2)^\top$ , how can you compute the probability that this observation comes from the \*second\* (i.e., Student's t copula) mixture component (given the outcome)? Provide the equation. [10pt]

### Question 3. [15pt]

On a sample of size  $n = 101$  you estimated

$$\bar{x} = (1 \ 4.9 \ 5.5 \ 0.7)^\top,$$

$$\mathcal{S} = \begin{pmatrix} 7 & 4 & -2 & -2 \\ 4 & 6 & -1 & 0 \\ -2 & -1 & c & 1 \\ -2 & 0 & 1 & 5 \end{pmatrix},$$

for some  $c > 0$ , where  $\bar{x}$  and  $\mathcal{S}$  denote the sample mean and covariance, respectively.

You formulate the joint hypothesis that the means of  $X_1$  and  $X_4$  are equal, \*and\* that the means of  $X_2$  and  $X_3$  are equal.

$\chi^2$ critical values for $\nu$ degrees of freedom	5%
$\nu = 1$	3.84
$\nu = 2$	5.99

$\nu = 3$	7.81
$\nu = 4$	9.49

A) At a 5% significance level, how large must  $c$  be for you to \*not reject\* your hypothesis using a  $\chi^2$  test? [15pt]

#### Question 4. [25pt]

A) In a PCA, explain why it is important to normalize the variables in terms of their means and standard deviations. [5pt]

B) Describe the main steps of one of the iterative (non-maximum-likelihood) techniques to estimate an orthogonal factor model. [10pt]

You estimated an orthogonal factor model and obtained the factor loading matrix

$$L = \begin{pmatrix} 2 & 3 \\ -1 & 1 \\ 1 & 4 \end{pmatrix}.$$

Your colleague says that you can multiply the factor loadings from the right by the matrix  $\mathcal{A}$  with

$$\mathcal{A} = \begin{pmatrix} 0.2 & -0.6 \\ 0.2 & 0.4 \end{pmatrix}$$

and obtain

$$\tilde{L} = L \cdot \mathcal{A} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 1 & 1 \end{pmatrix},$$

and that this model yields an orthogonal factor model that is indistinguishable from your model, and that therefore you can interpret the first factor as the average of the 1<sup>st</sup> and 3<sup>rd</sup> variable, and the second factor as the average of the 2<sup>nd</sup> and 3<sup>rd</sup> factor.

C) Comment whether your colleague is right or wrong. [10pt]

