

Exam Optimization of Business Processes

27 May 2014

This exam consists of 4 problems, each consisting of several questions.

All answers should be motivated, including calculations, formulas used, etc.

It is allowed to use 1 sheet of paper (or 2 sheets written on one side) with **hand-written** notes.

The minimal note is 1. All questions give the same number of points.

The use of a calculator and a dictionary are allowed.

A table with the normal distribution and a table with the Poisson distribution are attached.

1a. Let X be uniform on $[0, 1]$. Show that $\mathbb{E}X = 1/2$ and $\sigma^2(X) = 1/12$.

b. Let Y be uniform on $[a, b]$. Formulate Y in terms of X and use this and the answer of a to show that $\mathbb{E}X = 0.5(a + b)$ and $\sigma^2(X) = (b - a)^2/12$.

c. Consider an $M|G|1$ queue with mean interarrival time b and service time distribution Y . Compute the long-run average waiting time.

d. We split the customers up in 2 classes: those that have service time less than $\mathbb{E}Y$ and those that have a longer service time. Compute the long-run average waiting time for the priority policy that gives priority to short service times.

2. Consider a 1-out-of- k model with cold standby. All failed machines can be repaired in parallel. We assume that repair times and failure times are exponential with rates α and β .

a. Formulate the evolution of this system as a birth-death process.

b. Determine the probability that the system is down.

c. Compute this number for $k = 5$, $\alpha = 1$ and $\beta = 3$.

d. Suppose that the repair times are deterministic with value $1/\alpha$. Does this change the result? Motivate your answer.

- 3a. Formulate the call center shift scheduling problem that requires the service level to be attained in every interval.
- 3b. Give two reasons why this leads to overstaffing.
- 3c. Formulate the call center shift scheduling problem with an overall service level constraint.
- 3d. Formulate this as an integer programming problem with linear constraints.

- 4a. Consider a binomial distribution consisting of 100 experiments each with success probability 0.25. Give the parameters of the Poisson distribution and the normal distribution that best approximate this binomial distribution. Motivate your answer.
- 4b. Consider the Littlewood model with capacity 40, prices 1 and 3, and demand in both classes according to the binomial distribution just given. Use both approximations of question a to approximate the optimal level of capacity reserved for the expensive class.
- 4c. Formulate a dynamic programming recursion with 200 time periods that solves the same problem.

Table with values of $\mathbb{P}(X > k)$ with X a Poisson distributed random variable with mean μ

	values of μ									
values of k	1	2	3	4	5	6	7	8	9	10
0	0.632	0.865	0.950	0.982	0.993	0.998	0.999	1.000	1.000	1.000
1	0.264	0.594	0.801	0.908	0.960	0.983	0.993	0.997	0.999	1.000
2	0.080	0.323	0.577	0.762	0.875	0.938	0.970	0.986	0.994	0.997
3	0.019	0.143	0.353	0.567	0.735	0.849	0.918	0.958	0.979	0.990
4	0.004	0.053	0.185	0.371	0.560	0.715	0.827	0.900	0.945	0.971
5	0.001	0.017	0.084	0.215	0.384	0.554	0.699	0.809	0.884	0.933
6	0.000	0.005	0.034	0.111	0.238	0.394	0.550	0.687	0.793	0.870
7	0.000	0.001	0.012	0.051	0.133	0.256	0.401	0.547	0.676	0.780
8	0.000	0.000	0.004	0.021	0.068	0.153	0.271	0.407	0.544	0.667
9	0.000	0.000	0.001	0.008	0.032	0.084	0.170	0.283	0.413	0.542
10	0.000	0.000	0.000	0.003	0.014	0.043	0.099	0.184	0.294	0.417
11	0.000	0.000	0.000	0.001	0.005	0.020	0.053	0.112	0.197	0.303
12	0.000	0.000	0.000	0.000	0.002	0.009	0.027	0.064	0.124	0.208
13	0.000	0.000	0.000	0.000	0.001	0.004	0.013	0.034	0.074	0.136
14	0.000	0.000	0.000	0.000	0.000	0.001	0.006	0.017	0.041	0.083
15	0.000	0.000	0.000	0.000	0.000	0.001	0.002	0.008	0.022	0.049
16	0.000	0.000	0.000	0.000	0.000	0.000	0.001	0.004	0.011	0.027
17	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.002	0.005	0.014
18	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.001	0.002	0.007
19	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.001	0.003
20	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.002
21	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.001
22	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
23	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
24	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
25	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Table with values of $\mathbb{P}(0 < X < x + y)$ with X a random variable with a standard normal distribution

	values of y									
values of x	0	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
0	0.000	0.004	0.008	0.012	0.016	0.020	0.024	0.028	0.032	0.036
0.1	0.040	0.044	0.048	0.052	0.056	0.060	0.064	0.067	0.071	0.075
0.2	0.079	0.083	0.087	0.091	0.095	0.099	0.103	0.106	0.110	0.114
0.3	0.118	0.122	0.126	0.129	0.133	0.137	0.141	0.144	0.148	0.152
0.4	0.155	0.159	0.163	0.166	0.170	0.174	0.177	0.181	0.184	0.188
0.5	0.191	0.195	0.198	0.202	0.205	0.209	0.212	0.216	0.219	0.222
0.6	0.226	0.229	0.232	0.236	0.239	0.242	0.245	0.249	0.252	0.255
0.7	0.258	0.261	0.264	0.267	0.270	0.273	0.276	0.279	0.282	0.285
0.8	0.288	0.291	0.294	0.297	0.300	0.302	0.305	0.308	0.311	0.313
0.9	0.316	0.319	0.321	0.324	0.326	0.329	0.331	0.334	0.336	0.339
1	0.341	0.344	0.346	0.348	0.351	0.353	0.355	0.358	0.360	0.362
1.1	0.364	0.367	0.369	0.371	0.373	0.375	0.377	0.379	0.381	0.383
1.2	0.385	0.387	0.389	0.391	0.393	0.394	0.396	0.398	0.400	0.401
1.3	0.403	0.405	0.407	0.408	0.410	0.411	0.413	0.415	0.416	0.418
1.4	0.419	0.421	0.422	0.424	0.425	0.426	0.428	0.429	0.431	0.432
1.5	0.433	0.434	0.436	0.437	0.438	0.439	0.441	0.442	0.443	0.444
1.6	0.445	0.446	0.447	0.448	0.449	0.451	0.452	0.453	0.454	0.454
1.7	0.455	0.456	0.457	0.458	0.459	0.460	0.461	0.462	0.462	0.463
1.8	0.464	0.465	0.466	0.466	0.467	0.468	0.469	0.469	0.470	0.471
1.9	0.471	0.472	0.473	0.473	0.474	0.474	0.475	0.476	0.476	0.477
2	0.477	0.478	0.478	0.479	0.479	0.480	0.480	0.481	0.481	0.482
2.1	0.482	0.483	0.483	0.483	0.484	0.484	0.485	0.485	0.485	0.486
2.2	0.486	0.486	0.487	0.487	0.487	0.488	0.488	0.488	0.489	0.489
2.3	0.489	0.490	0.490	0.490	0.490	0.491	0.491	0.491	0.491	0.492
2.4	0.492	0.492	0.492	0.492	0.493	0.493	0.493	0.493	0.493	0.494
2.5	0.494	0.494	0.494	0.494	0.494	0.495	0.495	0.495	0.495	0.495
2.6	0.495	0.495	0.496	0.496	0.496	0.496	0.496	0.496	0.496	0.496
2.7	0.497	0.497	0.497	0.497	0.497	0.497	0.497	0.497	0.497	0.497
2.8	0.497	0.498	0.498	0.498	0.498	0.498	0.498	0.498	0.498	0.498
2.9	0.498	0.498	0.498	0.498	0.498	0.498	0.498	0.499	0.499	0.499
3	0.499	0.499	0.499	0.499	0.499	0.499	0.499	0.499	0.499	0.499