

HERTENTAMEN

Kansrekening en Statistiek – 4I

WEDNESDAY 31 AUGUST 2005

INSTRUCTIONS

- Write your name and studentnumber on *every* page
- Motivate all answers clearly
- Use of a calculator will *not* be necessary and is *not* allowed
- Use of Rice's book is allowed
- If you finish the exam early, hand in your work and leave *quietly*
- Do *not* attempt to cheat or confer with your fellow students

Points per question ($Grade = total/8 + 1$)

1(a):	5	2(a):	5	3(a):	5
1(b):	5	2(b):	5	3(b):	5
1(c):	5	2(c):	6	3(c):	5
1(d):	6	2(d):	5	3(d):	5
				3(e):	5
				3(f):	5

Good luck!

[DO NOT TURN THIS PAGE UNTIL SO INSTRUCTED]

PROBLEM 1

Let X in \mathbb{R} be continuously distributed with probability density function $f_X : \mathbb{R} \rightarrow \mathbb{R}$ of the form:

$$f_X(x) = \begin{cases} Cx^2(2-x)^2, & \text{if } 0 \leq x \leq 2, \\ 0, & \text{if } x < 0 \text{ or } x > 2. \end{cases}$$

a. (5 points)

Calculate the normalization constant C .

b. (5 points)

Show that $E(X) = 1$.

c. (5 points)

Show that the second moment $E(X^2)$ satisfies

$$E(X^2) = \frac{128}{105}C,$$

and, using your answer under part a., show that $\text{Var}(X) = 8/7$.

d. (6 points)

Give the distribution function $F_X : \mathbb{R} \rightarrow [0, 1]$ of X .

Based on your answer, demonstrate that the median of X equals 1.

PROBLEM 2

Let X, Y be two continuous random variables, jointly distributed according to a joint probability density function $f_{XY} : \mathbb{R}^2 \rightarrow \mathbb{R}$,

$$f_{XY}(x, y) = \begin{cases} Ce^{-\lambda(x+y)}, & \text{if } x \geq 0 \text{ and } y \geq 0, \\ Ce^{-\lambda(x-y)}, & \text{if } x \geq 0 \text{ and } y < 0, \\ Ce^{-\lambda(-x+y)}, & \text{if } x < 0 \text{ and } y \geq 0, \\ Ce^{\lambda(x+y)}, & \text{if } x < 0 \text{ and } y < 0. \end{cases}$$

depending on a parameter $\lambda > 0$.

a. (5 points)

Calculate the normalization constant C

(NB: your answer will be an expression involving λ).

b. (5 points)

Show that the marginal probability density function $f_X : \mathbb{R} \rightarrow \mathbb{R}$ for X has the following form:

$$f_X(x) = \begin{cases} \frac{\lambda}{2} e^{-\lambda x}, & \text{if } x \geq 0, \\ \frac{\lambda}{2} e^{\lambda x}, & \text{if } x < 0. \end{cases}$$

c. (6 points)

Show that the random variables X and Y are independent (motivating your answer by an equality of probability density functions).

What relation does the independence of X and Y provide between $\text{Var}(X+Y)$, $\text{Var}(X)$ and $\text{Var}(Y)$?

d. (5 points)

Show that:

$$\text{Var}(X + Y) = \frac{4}{\lambda^2}.$$

(NB: You may use that $\int_0^\infty u^n e^{-u} du = n!$.)

PROBLEM 3

Consider an experiment in which we measure X , a random variable with a distribution depending on an unknown parameter $\kappa > 0$. The probability density function $f_\kappa : \mathbb{R} \rightarrow \mathbb{R}$ for X is given by:

$$f_\kappa(x) = \begin{cases} 0, & \text{if } x < 0, \\ \frac{1}{\kappa\sqrt{\pi}} x^{-1/2} e^{-\frac{x}{\kappa^2}}, & \text{if } x \geq 0. \end{cases}$$

Assume that we have an *i.i.d.* sample X_1, X_2, \dots, X_n of repeated measurements of X . Parts *a.* and *b.* of this problem concern the moment estimator for κ .

a. (5 points)

By partial integration, show that:

$$\int_0^\infty x^{1/2} e^{-x/\kappa^2} dx = \frac{\kappa^2}{2} \int_0^\infty x^{-1/2} e^{-x/\kappa^2} dx.$$

Use this equality to show that $E(X) = \kappa^2/2$.

b. (5 points)

Give the first moment equation and derive the form of the moment estimator $\tilde{\kappa}$ for the parameter κ .

Indicate clearly in which step of your derivation the restriction $\kappa > 0$ plays a role.

Next, we turn to the maximum-likelihood estimator of κ . As it turns out, it is convenient to use, instead of κ itself, its square $\theta = \kappa^2$ as a parameter. This does not change the probability density functions in the model, only the parametrization. To obtain the maximum-likelihood estimator for κ , we shall solve the maximum-likelihood problem for θ first and then derive the maximum-likelihood estimator for κ .

c. (5 points)

What is the domain of the new parameter θ ?

Give the form for the probability density function $f_\theta : \mathbb{R} \rightarrow \mathbb{R}$ parametrized by the parameter θ .

Also give the log-likelihood $\ell(\theta)$ based on the *i.i.d.* sample X_1, \dots, X_n .

d. (5 points)

Write down the first derivative $\dot{\ell}(\theta)$ of $\ell(\theta)$ with respect to θ .

Solve the equation $\dot{\ell}(\theta) = 0$ to show that the maximum-likelihood estimator $\hat{\theta}$ for θ equals:

$$\hat{\theta}(X_1, \dots, X_n) = 2\bar{X}.$$

e. (5 points)

Based on part d., give the maximum-likelihood estimator $\hat{\kappa}$ for κ .

Explain your answer clearly.

(Hint: this does not require any calculation: extrema of a function are parametrization-invariant.)

f. (5 points)

Is the maximum-likelihood estimator $\hat{\theta}$ for θ biased or unbiased?

Can we conclude from this that the maximum-likelihood estimator $\hat{\kappa}$ for κ is biased/unbiased?