

Exam Measure Theory

October 24 2014, 8.45-11.30

Alle onderdelen tellen even zwaar mee.

Exercise 1. Let f be a function from $(\mathbb{R}, \mathcal{F}) \rightarrow (\mathbb{R}, \mathcal{B})$, where $\mathcal{F}$ is a sigma-algebra, and $\mathcal{B}$ denotes the Borel sigma-algebra.

(a) Does there exist a sigma-algebra $\mathcal{F}$ such that f is $\mathcal{F}/\mathcal{B}$ measurable if and only if f is a constant function? Explain your answer.

(b) Does there exist a sigma-algebra $\mathcal{F}$ such that f is $\mathcal{F}/\mathcal{B}$ measurable if and only if f is continuous? Explain your answer.

Exercise 2. For μ a pre-measure on a semi-ring $\mathcal{S}$ of subsets of a set X , we have defined the outer measure μ^* as follows:

$$\mu^*(A) := \inf \left\{ \sum_{j=1}^{\infty} \mu(S_j) : S_j \in \mathcal{S}, \bigcup_{j=1}^{\infty} S_j \supset A \right\}.$$

(a) Show that in case $X = \mathbb{R}$, the collection $\mathcal{S}$ of intervals of the form $[a, b)$, $a, b \in \mathbb{R}$ is a semi-ring.

(b) We have shown in class that Lebesgue measure λ is a pre-measure on the semi-ring in (a). Show that in this case,

$$\mu^*(A) = \inf \left\{ \sum_{j=1}^{\infty} \mu(S_j) : S_j \text{ is open}, \bigcup_{j=1}^{\infty} S_j \supset A \right\}.$$

You can use the fact that open sets in $\mathbb{R}$ are countable unions of open intervals.

Exercise 3. Let $x \in [0, 1]$ and write x in binary representation as

$$x = 0.a_1(x)a_2(x)a_3(x)\dots$$

where $a_n(x) \in \{0, 1\}$. By this we mean that

$$x = \sum_{n=1}^{\infty} \frac{a_n(x)}{2^n}.$$

(Some x have two such representations. In those cases we choose the one for which $a_n(x) = 1$ for all large enough n .) Let $f : [0, 1] \rightarrow [0, 1]$ be defined by

$$f(x) = 0.a_2(x)a_3(x)a_4(x)\dots = \sum_{n=1}^{\infty} \frac{a_{n+1}(x)}{2^n}.$$

- (a) Show that for all n , $a_n : [0, 1] \rightarrow [0, 1]$ is $\mathcal{B}/\mathcal{B}$ measurable.
(b) Let, for $k = 1, 2, \dots$

$$f_k(x) = \sum_{n=1}^k \frac{a_{n+1}(x)}{2^n}.$$

Show that f_k is measurable and use this to show that f is measurable.

- (c) Show that f is piecewise continuous.
(d) Use (c) to give a second proof of the measurability of f .

Exercise 4. Let $A_1, A_2, \dots$ be elements of $\mathcal{B}$ such that $\lambda(A_m \cap A_n) = 0$ for all $m \neq n$. Let $B_n = A_n \cap A_1^c \cap A_2^c \cap \dots \cap A_{n-1}^c$.

- (a) Show that

$$\lambda\left(\bigcup_{n=1}^{\infty} A_n\right) = \lambda\left(\bigcup_{n=1}^{\infty} B_n\right) = \sum_{n=1}^{\infty} \lambda(B_n).$$

- (b) Show that

$$A_n \triangle B_n \subset \bigcup_{m=1}^n (A_m \cap A_n).$$

(Here $\triangle$ denotes symmetric difference.)

- (c) Show that

$$\lambda\left(\bigcup_{n=1}^{\infty} A_n\right) = \sum_{n=1}^{\infty} \lambda(A_n).$$

Exercise 5. Let $\mathbb{N} = \{1, 2, 3, \dots\}$ and define the sets $A_k \subset \mathbb{N}$ by

$$A_k = \{k, 2k, 3k, \dots\},$$

for $k = 1, 2, \dots$. We denote by $\mathcal{H}$ the collection $\{A_1, A_2, \dots\} \cup \emptyset$.

- (a) Show that $\sigma(\mathcal{H})$ (the sigma-algebra generated by $\mathcal{H}$) is equal to $\mathcal{P}(\mathbb{N})$ (the power set of $\mathbb{N}$).
(b) Suppose that μ and ν are finite measure on $(\mathbb{N}, \mathcal{P}(\mathbb{N}))$ such that $\mu(H) = \nu(H)$ for all $H \in \mathcal{H}$. Show that $\mu = \nu$.