

**Use of a basic calculator is allowed. Graphical calculators and mobile phones are not allowed. This exam consists of 4 questions (27 points).**

**Please write all answers in English. Grade =  $\frac{total+3}{3}$ .**

### GOOD LUCK!

#### Question 1 [6 points]

Are the following statements correct/sensible? Motivate your answer by a short argument.

- [2 points] The Shapiro-Wilk test is for testing a composite null hypothesis.
- [2 points]  $M$ -estimators with bounded  $\psi$ -function are robust.
- [2 points] A two sample  $QQ$ -plot is the same as a scatter plot for paired data.

#### Question 2 [6 points]

Let  $X_1, \dots, X_{100}$  be independent and identically distributed random variables with unknown distribution  $P$ . Suppose we want to test  $H_0 : P = P_0$ , for a known  $P_0$ , using the  $\chi^2$  goodness-of-fit test.

- [3 points] The test statistic

$$X^2 = \sum_{i=1}^k \frac{(N_i - np_i)^2}{np_i}$$

has approximately a  $\chi^2_{k-1}$ -distribution under  $H_0$ . Describe the rule of thumb that needs to be satisfied for this approximation to be reliable.

- [3 points] Suppose we are given intervals  $I_1, \dots, I_k$  that do not fulfill the rule of thumb for the given sample size. In such a situation we can still use  $X^2$  as test statistic. However, we cannot rely on its approximate  $\chi^2$ -distribution. Therefore, we use a bootstrap test. Describe the steps that are made in a bootstrap test for the given null hypothesis using  $X^2$  as test statistic.

#### Question 3 [7 points]

In Figure 1 a histogram, boxplot and several  $QQ$ -plots of a data set  $x$  are presented.

- [2 point] Which of the four location scale families do you think is most appropriate for these data? Explain your answer.




Figure 1: Histogram, boxplot and QQ-plots against the  $N(0,1)$ , uniform $[0,1]$ , exponential(1) and the standard  $\chi^2_8$  distributions of a data set.

- b. [2 points] The  $\alpha$ -trimmed mean of these data was computed for  $\alpha = 0, 0.1, 0.2, 0.3, 0.4, 0.5$ . The values of these 6 trimmed means are, in arbitrary order: 0.69, 0.62, 0.81, 0.63, 0.74, 1.02. Which of these values is the 0.1 trimmed mean? Motivate your answer clearly!
- c. [3 points] Using the QQ-plot you have selected under part (a) determine the location  $a$  and scale  $b$  approximately. You may use that the sample variance equals 1.32. (You may use that the expectation and variance of the  $\chi^2_k$ -distribution are  $k$  and  $2k$ , respectively.)

#### Question 4 [8 points]

Let  $X_1, \dots, X_n$  be independent and identically distributed random variables with unknown distribution  $P$ . Suppose that the sample variance  $T_n(X_1, \dots, X_n) = S_X^2$  is used to estimate the variance of  $P$ . To determine the accuracy of this estimator, its standard deviation is estimated by means of the empirical bootstrap.

- a. [4 points] Describe the steps of the empirical bootstrap scheme that you would use to find the bootstrap estimate of the standard deviation of  $T_n$ .
- b. [2 points] Describe shortly which two errors are (necessarily) made in this bootstrap procedure.
- c. [2 points] Which of the two errors in part (b) can be made arbitrarily small? What do you have to change in the procedure under (a) to make this error smaller?